

Talk given at XXI International Meeting on Fundamental Physics - Physics at HERA, Miraflores de la Sierra, Madrid, Spain, 9-13 May 1993.

QED Radiative Corrections to Deep Inelastic Scattering at HERA

A. A. Akhundov

*DESY - Institut für Hochenergiephysik Zeuthen,
Platanenallee 6, D-15738 Zeuthen, Germany*

ABSTRACT

The numerical study of QED radiative corrections in the low x region for the H1 experiment at HERA was performed. The analytical program TERAD91 was used to obtain the values of the radiative correction factor for the first measurement of the proton structure function $F_2(x, Q^2)$ in the range $x = 10^{-2} - 10^{-4}$ and $5 < Q^2 < 80 \text{ GeV}^2$.

In order to extract from experimental data the inclusive cross section of deep inelastic ep -scattering

$$e(k_1) + p(p_1) \rightarrow e(k_2) + X(p_2) \quad (1)$$

in one-boson-exchange approximation one must take into account contributions from higher order QED¹ and electroweak processes. For example the bremsstrahlung process:

$$e(k_1) + p(p_1) \rightarrow e(k_2) + X(p_2) + \gamma(k), \quad (2)$$

and also the elastic radiative tail:

$$e(k_1) + p(p_1) \rightarrow e(k_2) + p(p_2) + \gamma(k). \quad (3)$$

contribute to the observed cross section of the reaction (1).

In the long period from the SLAC experiments² till the operation of HERA, the experiments with neutral current deep inelastic ep -scattering had a relatively simple structure, and the measurements completely rested on the registration of energy and angle of the scattered electron. The new detector generation, the HERA detectors H1³ and ZEUS,⁴ allows to measure both the scattered electron and hadronic final state.

For the physical analysis of the ep -collisions at HERA one can use not only the familiar electron variables

$$Q_e^2 = -(k_1 - k_2)^2, \quad y_e = p_1(k_1 - k_2)/p_1 k_1, \quad x_e = Q_e^2/(sy_e), \quad (4)$$

where

$$s = (k_1 + p_1)^2 = 4E_e E_p, \quad (5)$$

but also the kinematical variables from the hadron measurement

$$Q_h^2 = -(p_2 - p_1)^2, \quad y_h = p_1(p_2 - p_1)/p_1 k_1, \quad x_h = Q_h^2/(s y_h), \quad (6)$$

or some mixture of both. Here E_e and E_p are the energies of incident electron and proton.

For different choices of variables there are differences in the predictions for the contribution of the higher order processes.

The theoretical investigations of these radiative effects, or radiative corrections (RC), for experiments at HERA were summarized in Proceedings of the Workshop "Physics at HERA" .⁵ Different programs to calculate the radiative corrections have been cross-checked .⁶

The analysis of neutral current deep inelastic scattering data in the H1 experiment at HERA in 1992⁷ was based on two independent approaches:

- i) a "detector oriented" approach (analysis I)
- ii) a "physics oriented" approach (analysis II)

Within the analysis I the kinematics of the inclusive deep inelastic scattering process (1) were determined from the energy E'_e of the scattered electron, and its polar angle θ_e , measured relative to the proton beam direction

$$Q_e^2 = 4E_e E'_e \cos^2(\theta_e/2), \quad (7)$$

$$y_e = 1 - (E'_e/E_e) \sin^2(\theta_e/2). \quad (8)$$

The cross sections, acceptances of detector, radiative corrections and resolution effects were obtained in $(\sqrt{E'_e}, \theta_e)$ bins. The cross sections, that have been measured using this binning, were transformed into cross sections in (x_e, Q_e^2) -bins.

In the analysis II the kinematics of the process (1) were obtained from a measurement of the electron variables E'_e , θ_e and scaling variable y_h from the hadrons. The hadronic variable y_h (6) can be determined using the Jacquet-Blondel method⁸

$$y_h = \sum_{hadrons} \frac{E_h - p_{z,h}}{2E_e}, \quad (9)$$

where E_h is the energy of a hadron and $p_{z,h}$ its momentum component along the incident proton direction.

The momentum transfer Q^2 was determined from the electron variables θ_e and E'_e as in (7) and Bjorken x by using a mixture of the electron and hadron measurements⁹

$$x_m = Q_e^2/(s y_h). \quad (10)$$

In the analysis II cross sections, radiative corrections and efficiencies were calculated in (x_m, Q_e^2) -bins.

A further difference between the two methods of the analysis of the data was the use of two different types of programs — the Monte Carlo event generators HERACLES¹⁰ in the analysis I and the analytical program TERAD91¹¹ in the analysis II — for the calculations of the radiative correction factor $\delta(x, Q^2)$. The radiative correction factor $\delta(x, Q^2)$ is defined by the equation

$$\delta(x, Q^2) = \frac{d^2\sigma^{\text{theor}}/dxdQ^2}{d^2\sigma^{\text{Born}}/dxdQ^2} - 1, \quad (11)$$

where $d^2\sigma^{\text{theor}}/dxdQ^2$ is the theoretical approximation to the measured cross section $d^2\sigma^{\text{meas}}/dxdQ^2$, which contains the contributions from higher order electroweak processes, and $d^2\sigma^{\text{Born}}/dxdQ^2$ is the Born cross section of the process (1).

In this note we describe the numerical calculations of the RC factor $\delta(x, Q^2)$ for the "Zeuthen analysis" of the proton structure function $F_2(x, Q^2)$ in the range $x = 10^{-2} - 10^{-4}$ and $5 < Q^2 < 80 \text{ GeV}^2$ in the H1 experiment at HERA.⁷

The Monte Carlo HERACLES is suited for the calculation of RC to the differential cross sections in terms of the electron variables because the program calculates cross sections and generates events in terms of the variables x_e and Q_e^2 . Therefore most of the generated radiative events in HERACLES have low Q_h^2 and the calculation of the radiative corrections in the terms of other variables (hadron, mixed) can be performed with less efficiency only. For this reason in the analysis II was used the analytical program TERAD91.

TERAD91 - a program package to calculate radiative corrections for deep inelastic ep -scattering - was created in 1991 during Workshop "Physics at HERA" and accumulates about 15 years experience in the field of calculation of the radiative corrections for deep inelastic lN -scattering.¹⁴ The program TERAD91 can be applied to calculate the RC factor of the order $\mathcal{O}(\alpha)$ to the measured differential cross section, $d^2\sigma^{\text{meas}}/dxdQ^2$, after the acceptance and smearing corrections.

The program package TERAD91 contains one part, TERAD, which is based on the model-independent treatment¹⁵ of the leptonic QED corrections for neutral current ep -scattering, including bremsstrahlung process (2) and QED vertex corrections. TERAD does not necessarily rely on the quark parton model and uses the phenomenological structure functions F_1, F_2 and F_3 , which are defined in the Born approximation.

The program TERAD91 can be used for the model-independent calculations of QED corrections for neutral current deep inelastic ep -scattering in the terms of several variables - electron, hadron, mixed variables¹⁶- and in the new version¹⁷ also in terms of the Jacquet-Blondel variables.¹⁸

In the low Q^2 region the Born cross section of the deep inelastic scattering (1)

is determined by two structure function F_2 and $2xF_1 = F_2/(1 + R)$:

$$\frac{d^2\sigma}{dx dQ^2} = \frac{2\pi\alpha^2}{Q^4 x} \left(2(1 - y) + \frac{y^2}{1 + R} \right) F_2(x, Q^2). \quad (12)$$

In order to calculate the RC one has to use a realistic parametrization of the structure functions $F_i(x, Q^2)$ over the full range of x and Q^2 . We have used several different parameterizations of the deep inelastic proton structure functions, including MRS D- and MRS D0 parameterizations.¹³ For region $Q^2 < 5 \text{ GeV}^2$ the parameterizations were taken at $Q^2 = 5 \text{ GeV}^2$ and multiplied by factor¹⁹ $(1 - \exp(-aQ^2))$, $a = 3.37 \text{ GeV}^{-2}$. In our calculations we have assumed $R = 0$.

In the "Zeuthen analysis" the following x, Q^2 -binning was used. There were four Q^2 -bins:

$$Q^2 = 5 - 10, 10 - 20, 20 - 40, 40 - 80 \text{ GeV}^2 \quad (13)$$

and four bins per x -decade:

$$\log x = -4.0, -3.75, -3.50, -3.25, \dots, -1.0. \quad (14)$$

The kinematical region of the experimental measurement was defined by:

$$157.5^\circ < \theta_e < 172.5^\circ, \quad 0.05 < y_e < 0.60 \quad (15)$$

for electron variables and

$$157.5 < \theta_e < 172.5^\circ, \quad y_h < 0.50, \quad \theta_{jet} > 10^\circ \quad (16)$$

for mixed variables.

Figs.1,2 show the results for the radiative corrections in the terms of the mixed variables x_m, Q_e^2 . The RC in the terms of the mixed variables, $\delta(x_m, Q_e^2)$, are not small and can be reach 7 % at $x = 10^{-2}$.

The "Zeuthen analysis" of the deep inelastic scattering has included also the analysis of the experimental data in the terms of the electron variables x_e, Q_e^2 corresponding to the binning (13), (14). Fig.3,4 show the results of the calculations of the RC in the bins x_e, Q_e^2 , that have been obtained with the program TERAD91.

The corrections determined by the electron variables, $\delta(x_e, Q_e^2)$, are known to be large at low x and dominated by hard photon emission from the incoming electron. From the comparison of Fig.3 and Fig.4 it is seen that the sensitivity of the RC to the choice of the structure functions is not small.

Because the collinear radiation of photons in the final state of the reaction (2) is not resolved in the electromagnetic calorimeter, one must calculate the RC in terms of the calorimetric variables including the total energy of the electromagnetic cluster

$$E_{vis} = E'_e + E_\gamma. \quad (17)$$

In the program TERAD91 the possibility to calculate the RC in terms of calorimetric variables, E_{vis}, θ_e , has not been included yet, but with the help of the leading log calculations¹² one can estimate the size of the influence of the calorimetric measurement on the radiative corrections.

The calorimetric measurement cannot distinguish between a single electron and an electron with accompanying photons. Therefore the final state radiation has experimentally no effect on the measurement of the kinematics of deep inelastic scattering and can be absorbed in the definition of the final electron energy of the process (1).

This leads to the following way to take into account the calorimetric measurement. We can calculate the contribution of the final state radiation in the leading logarithmic approximation and subtract from the total RC, that we obtain with the program TERAD91. The more detailed investigation of the calorimetry problem composes the subject of a separate publication.

Acknowledgements

I would like to thank D. Bardin and M. Klein for numerous discussion. It is a great pleasure to thank the DESY–IfH Zeuthen for the opportunities to work at DESY.

I would like to acknowledge F. Barreiro for the kind invitation to participate in this Meeting and P. Söding for the financial support.

References

1. L. W. Mo and Y. S. Tsai, *Rev. Mod. Phys.* **41** (1969) 205;
Y. S. Tsai, preprint SLAC-Pub-848 (1971).
2. E. D. Bloom et al., *Phys. Rev. Lett.* **23** (1969) 930.
M. Breidenbach et al., *Phys. Rev. Lett.* **23** (1969) 935.
3. I. Abt et al., *The H1 Detector at HERA*, DESY preprint 93-103 (1993).
4. M. Derrick et al., *Phys. Lett.* **B293** (1992) 465.
5. *Proceedings of the Workshop "Physics at HERA"*, W. Buchmüller and G. Ingelman (eds.), Hamburg (1991)
6. H. Spiesberger et al. , in *Proceedings of the Workshop "Physics at HERA"*, ed. W. Buchmüller and G. Ingelman, Hamburg, (1991)p. 798.
7. A. De Roeck, *Deep Inelastic Scattering at Low-x. Results from the H1 Experiment*, DESY preprint 93-087 (1993);
I. Abt et al., *Measurement of the Proton Structure Function $F_2(x, Q^2)$ in the Low x Region at HERA*, DESY preprint 93-117 (1993).
8. A. Blondel and F. Jacquet, in *Proceedings of the Study of an ep Facility for Europe*, ed. U. Amaldi, DESY preprint 79-48 (1979) 391.
9. M. Klein, in *Proceedings of 4th San Miniato Topical Seminar*, ed. P. Giusti et al, Singapore (1990)p. 31
10. A. Kwiatkowski, H. Spiesberger, and H.-J. Möhring, *Comp. Phys. Commun.* **69** (1992) 155.

11. A. Akhundov et al., in *Proceedings of the Workshop "Physics at HERA"*, ed. W. Buchmüller and G. Ingelman, Hamburg, (1991) p. 1285.
12. J. Blümlein, *Z. Phys.* **C47** (1990) 89; *Phys. Lett.* **B271** (1991).
13. A. D. Martin, W. J. Stirling, R. G. Roberts, *Phys. Rev.* **D47** (1993) 867.
14. A. A. Akhundov, D. Yu. Bardin, N. M. Shumeiko, JINR Dubna preprint E2-10147 and E2-10205 (1976);
A. Akhundov, D. Bardin, N. Shumeiko, *Sov. J. Nucl. Phys.* **26** (1977) 660;
N. Shumeiko, *Sov. J. Nucl. Phys.* **29** (1979) 807;
A. Akhundov, D. Bardin, W. Lohmann, JINR Dubna preprint E2-86-104 (1986) and program **TERAD86**;
A. A. Akhundov, D. Yu. Bardin, N. M. Shumeiko, *Sov. J. Nucl. Phys.* **44** (1986) 988;
D. Bardin, C. Burdik, P. Christova and T. Riemann, *Z. Phys.* **C42** (1989) 679;
D. Bardin, C. Burdik, P. Christova and T. Riemann, *Z. Phys.* **C44** (1989) 149;
A.A. Akhundov et al., *Z. Phys.* **C45** (1990) 645.
15. A. Akhundov, D. Bardin, L. Kalinovskaya and T. Riemann, in preparation.
16. D. Bardin, A. Akhundov, L. Kalinovskaya and T. Riemann, in *Proceedings of the Zeuthen Workshop on Elementary Particle Theory – Deep Inelastic Scattering, Teupitz, 1992*, ed. J. Blümlein and T. Riemann, *Nucl. Phys. B* (Proc. Suppl.) **29A** (1992) 209
17. A. Akhundov et al., DESY preprint, in preparation.
18. A. Akhundov, D. Bardin, L. Kalinovskaya and T. Riemann, *Phys. Lett.* **301B** 447.
19. N. Yu. Volkonsky and L. V. Prokhorov, *Sov. J. Exp. Theor. Phys. Lett.* **21** (1975) 389.




Fig. 1. Radiative correction factor $\delta(x_m, Q_e^2)$ in (x, Q^2) -bins for MRS D- parametrization.




Fig. 2. Radiative correction factor $\delta(x_m, Q_e^2)$ in (x, Q^2) -bins for MRS D0 parametrization.




Fig. 3. Radiative correction factor $\delta(x_e, Q_e^2)$ in (x, Q^2) -bins for MRS D- parametrization.




Fig. 4. Radiative correction factor $\delta(x_e, Q_e^2)$ in (x, Q^2) -bins for MRS D0 parametrization.